

## Soil Quality Indicators: pH

USDA Natural Resources Conservation Service

January 1998

### What is pH?

Soil pH is a measure of the acidity or alkalinity in the soil. It is also called soil reaction.

The most common classes of soil pH are:

Extremely acid	3.5 – 4.4
Very strongly acid	4.5 – 5.0
Strongly acid	5.1 – 5.5
Moderately acid	5.6 – 6.0
Slightly acid	6.1 – 6.5
Neutral	6.6 – 7.3
Slightly alkaline	7.4 – 7.8
Moderately alkaline	7.9 – 8.4
Strongly alkaline	8.5 – 9.0



### What is the significance of pH?

#### Availability of Nutrients

Soil pH influences the solubility of nutrients. It also affects the activity of micro-organisms responsible for breaking down organic matter and most chemical transformations in the soil. Soil pH thus affects the availability of several plant nutrients.

A pH range of 6 to 7 is generally most favorable for plant growth because most plant nutrients are readily available

in this range. However, some plants have soil pH requirements above or below this range.

Soils that have a pH below 5.5 generally have a low availability of calcium, magnesium, and phosphorus. At these low pH's, the solubility of aluminum, iron, and boron is high; and low for molybdenum.

At pH 7.8 or more, calcium and magnesium are abundant. Molybdenum is also available if it is present in the soil minerals. High pH soils may have an inadequate availability of iron, manganese, copper, zinc, and especially of phosphorus and boron.

#### Micro-organisms

Soil pH affects many micro-organisms. The type and population densities change with pH. A pH of 6.6 to 7.3 is favorable for microbial activities that contribute to the availability of nitrogen, sulfur, and phosphorus in soils.

#### Pesticide Interaction

Most pesticides are labeled for specific soil conditions. If soils have a pH outside the allowed range, the pesticides may become ineffective, changed to an undesirable form, or may not degrade as expected, which results in problems for the next crop period.

#### Mobility of heavy metals

Many heavy metals become more water soluble under acid conditions and can move downward with water through the soil, and in some cases move to aquifers, surface streams, or lakes.

#### Corrosivity

Soil pH is one of several properties used as a general indicator of soil corrosivity. Generally, soils that are either highly alkaline or highly acid are likely to be corrosive to steel. Soils that have pH of 5.5 or lower are likely to be highly corrosive to concrete.

### What controls soil pH?

The acidity or alkalinity in soils have several different sources. In natural systems, the pH is affected by the mineralogy, climate, and weathering. Management of soils

often alters the natural pH because of acid-forming nitrogen fertilizers, or removal of bases (potassium, calcium, and magnesium). Soils that have sulfur-forming minerals can produce very acid soil conditions when they are exposed to air. These conditions often occur in tidal flats or near recent mining activity where the soil is drained.

The pH of a soil should always be tested before making management decisions that depend on the soil pH.

## How is pH measured?

A variety of kits and devices are available to determine the pH in the field. The methods include:

- dyes
- paper strips
- glass electrodes.

Soil pH can change during the year. It depends on temperature and moisture conditions, and can vary to as much as a whole pH unit during the growing season. Since pH is a measure of the hydrogen ion activity [H<sup>+</sup>], many different chemical reactions can affect it. Temperature changes the chemical activity, so most measurements of pH include a temperature correction to a standard temperature of 25 degrees C (77°F). The soil pH generally is recorded as a range in values for the soil depth selected.



(Prepared by the National Soil Survey Center in cooperation with the Soil Quality Institute, NRCS, USDA, and the National Soil Tilth Laboratory, Agricultural Research Service, USDA).

The U. S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

## How is soil pH modified?

A soil pH below about 5.6 is considered low for most crops. Generally, the ideal pH range is between 6.0 and 7.0. Liming is a common method to increase the pH. It involves adding finely ground limestone to the soil. The reaction rate for limestone increases when soil temperatures are warm and soil moisture is high. If the limestone is more finely ground, the reaction is faster.

The amount of limestone to apply depends on the amount of organic matter and clay as well as the pH. Fertility testing laboratories that have local experience make this determination.

A soil pH that is more than about 8.0 is considered high for most crops. Soils that have a pH in this range are often also calcareous.

Calcareous soils have a high content of calcium carbonate. The pH of these soils does not change until most of the calcium carbonate is removed. Acids that are added to the soil dissolve the carbonates and lower the soil pH. Treatments with acid generally are uneconomical for soils that have a content of calcium carbonate of more than about 5%. Because phosphorus, iron, copper, and zinc are less available to plants in calcareous soils, nutrient deficiencies are often apparent. Applications of these nutrients are commonly more efficient than trying to lower the pH.

When the soil pH is above 8.6, sodium often is present. These soils generally do not have gypsum or calcium carbonates, at least not in the affected soil horizons. Addition of gypsum followed by leaching using irrigation is a common reclamation practice. However, salts flushed into drainage water may contaminate downstream waters and soils.

The application of anhydrous ammonia as a nitrogen fertilizer contributes to lowering the soil pH. In some parts of the country, applications of ammonia lower the surface soil pH from ranges of 6.6 to 7.3 to below 5.6. This reduction can be easily overlooked in areas of no-till cropping unless the pH is measured in the upper 2 inches.

Chemical amendments that contain sulfur generally form an acid, which lowers the soil pH.

**Visit our Web site:**  
<http://soils.usda.gov>